

References for TMD and Occlusal Factors

- Mohlin BO, Derweduwen K, Pilley R, Kingdon A, Shaw WC, Kenealy P.**
Malocclusion and temporomandibular disorder: a comparison of adolescents with moderate to severe dysfunction with those without signs and symptoms of temporomandibular disorder and their further development to 30 years of age. Angle Orthod. 2004 Jun;74(3):319-27.
- Tuerlings V, Limme M.**
The prevalence of temporomandibular joint dysfunction in the mixed dentition. Eur J Orthod. 2004 Jun;26(3):311-20.
- Gesch D, Bernhardt O, Kirbschus A.**
Association of malocclusion and functional occlusion with temporomandibular disorders (TMD) in adults: a systematic review of population-based studies. Quintessence Int. 2004 Mar;35(3):211-21.
- Fujita Y, Motegi E, Nomura M, Kawamura S, Yamaguchi D, Yamaguchi H.**
Oral habits of temporomandibular disorder patients with malocclusion. Bull Tokyo Dent Coll. 2003 Nov;44(4):201-7.
- Barker DK.**
Occlusal interferences and temporomandibular dysfunction. Gen Dent. 2004 Jan-Feb;52(1):56-61; quiz 62.
- Ali IM, Yamada K, Alkhamrah B, Vergara R, Hanada K.**
Relationship between occlusal curvatures and mandibular deviation in orthodontic patients with temporomandibular disorders. J Oral Rehabil. 2003 Nov;30(11):1095-103.
- Henrikson T, Nilner M.**
Temporomandibular disorders, occlusion and orthodontic treatment. J Orthod. 2003 Jun;30(2):129-37; discussion 127.
- Egermark I, Magnusson T, Carlsson GE.**
A 20-year follow-up of signs and symptoms of temporomandibular disorders and malocclusions in subjects with and without orthodontic treatment in childhood. Angle Orthod. 2003 Apr;73(2):109-15.
- Mohlin B, Kurol J.**
To what extent do deviations from an ideal occlusion constitute a health risk? Swed Dent J. 2003;27(1):1-10.
- Kusayama M, Motohashi N, Kuroda T.**
Relationship between transverse dental anomalies and skeletal asymmetry. Am J Orthod Dentofacial Orthop. 2003 Mar;123(3):329-37.
- Mohlin B, al-Saadi E, Andrup L, Ekblom K.**
Orthodontics in 12-year old children. Demand, treatment motivating factors and treatment decisions. Swed Dent J. 2002;26(2):89-98.
- Carlsson GE, Egermark I, Magnusson T.**
Predictors of signs and symptoms of temporomandibular disorders: a 20-year follow-up study from childhood to adulthood. Acta Odontol Scand. 2002 Jun;60(3):180-5.

references page 2.

Marzooq AA, Yatabe M, Ai M.

What types of occlusal factors play a role in temporomandibular disorders...? A literature review. J Med Dent Sci. 1999 Sep;46(3):111-6.

Celic R, Jerolimov V.

Association of horizontal and vertical overlap with prevalence of temporomandibular disorders. J Oral Rehabil. 2002 Jun;29(6):588-93.

Thilander B, Rubio G, Pena L, de Mayorga C.

Prevalence of temporomandibular dysfunction and its association with malocclusion in children and adolescents: an epidemiologic study related to specified stages of dental development. Angle Orthod. 2002 Apr;72(2):146-54.

Celic R, Jerolimov V, Panduric J.

A study of the influence of occlusal factors and parafunctional habits on the prevalence of signs and symptoms of TMD. Int J Prosthodont. 2002 Jan-Feb;15(1):43-8.

John MT, Hirsch C, Drangsholt MT, Mancl LA, Setz JM.

Overbite and overjet are not related to self-report of temporomandibular disorder symptoms. J Dent Res. 2002 Mar;81(3):164-9.

Tullberg M, Tsarapatsani P, Huggare J, Kopp S.

Long-term follow-up of early treatment of unilateral forced posterior cross-bite with regard to temporomandibular disorders and associated symptoms. Acta Odontol Scand. 2001 Oct;59(5):280-4.

Sonnesen L, Bakke M, Solow B.

Temporomandibular disorders in relation to craniofacial dimensions, head posture and bite force in children selected for orthodontic treatment. Eur J Orthod. 2001 Apr;23(2):179-92.

Gaudet EL Jr, Brown DT.

Temporomandibular disorder treatment outcomes: first report of a large-scale prospective clinical study. Cranio. 2000 Jan;18(1):9-22.

Zhang ZK, Ma XC, Gao S, Gu ZY, Fu KY.

Studies on contributing factors in temporomandibular disorders. Chin J Dent Res. 1999 Dec;2(3-4):7-20.

Seligman DA, Pullinger AG.

Analysis of occlusal variables, dental attrition, and age for distinguishing healthy controls from female patients with intracapsular temporomandibular disorders. J Prosthet Dent. 2000 Jan;83(1):76-82.

Pullinger AG, Seligman DA.

Quantification and validation of predictive values of occlusal variables in temporomandibular disorders using a multifactorial analysis. J Prosthet Dent. 2000 Jan;83(1):66-75.

references page 3.

Liu JK, Tsai MY.

Association of functional malocclusion with temporomandibular disorders in orthodontic patients prior to treatment. *Funct Orthod.* 1998 Jul-Sep;15(3):17-20.

Fushima K, Inui M, Sato S.

Dental asymmetry in temporomandibular disorders. *J Oral Rehabil.* 1999 Sep;26(9):752-6.

Kahn J, Tallents RH, Katzberg RW, Ross ME, Murphy WC.

Prevalence of dental occlusal variables and intraarticular temporomandibular disorders: molar relationship, lateral guidance, and nonworking side contacts. *J Prosthet Dent.* 1999 Oct;82(4):410-5.

Crawford SD.

Condylar axis position, as determined by the occlusion and measured by the CPI instrument, and signs and symptoms of temporomandibular dysfunction. *Angle Orthod.* 1999 Apr;69(2):103-15; discussion 115-6.

Sonnesen L, Bakke M, Solow B.

Malocclusion traits and symptoms and signs of temporomandibular disorders in children with severe malocclusion. *Eur J Orthod.* 1998 Oct;20(5):543-59.

Liu JK, Tsai MY.

Relationship between morphologic malocclusion and temporomandibular disorders in orthodontic patients prior to treatment. *Funct Orthod.* 1997 Nov-Dec;14(5):13-6.

Orbach S.

TMD issues in patient with skeletal deformity of jaws. *N Y State Dent J.* 1998 Mar;64(3):48-54.

Morrant DG, Taylor GS.

The prevalence of temporomandibular disorder in patients referred for orthodontic assessment. *Br J Orthod.* 1996 Aug;23(3):261-5.

Becker IM.

Occlusion as a causative factor in TMD. Scientific basis to occlusal therapy. *N Y State Dent J.* 1995 Nov;61(9):54-7.

Greene CS.

Etiology of temporomandibular disorders. *Semin Orthod.* 1995 Dec;1(4):222-8.

Raustia AM, Pirttiniemi PM, Pyhtinen J.

Correlation of occlusal factors and condyle position asymmetry with signs and symptoms of temporomandibular disorders in young adults. *Cranio.* 1995 Jul;13(3):152-6.

Pullinger AG, Seligman DA, Gornbein JA.

A multiple logistic regression analysis of the risk and relative odds of temporomandibular disorders as a function of common occlusal features. *J Dent Res.* 1993 Jun;72(6):968-79.

Tanne K, Tanaka E, Sakuda M.

Association between malocclusion and temporomandibular disorders in orthodontic patients before treatment. *J Orofac Pain.* 1993 Spring;7(2):156-62.

references page 4.

Hirata RH, Heft MW, Hernandez B, King GJ.

Longitudinal study of signs of temporomandibular disorders (TMD) in orthodontically treated and nontreated groups. Am J Orthod Dentofacial Orthop. 1992 Jan;101(1):35-40.

Kampe T, Hannerz H.

Five-year longitudinal study of adolescents with intact and restored dentitions: signs and symptoms of temporomandibular dysfunction and functional recordings. J Oral Rehabil. 1991 Sep;18(5):387-98.